An In-Depth Analysis of the Analgesic, Antimicrobial, and Anti-Inflammatory Effects of Essential Oils

Sidra Zubair[1], Fatima Qamar[2], Anum Liaqat[3],Quratulain Waseem[4],Shumaila Qadir[4]

[1]Department of Pharmacology, Faculty of Pharmacy, Jinnah University for Women, Karachi, Pakistan
[2]Department of Pharmaceutical Chemistry, Jinnah University for Women, Karachi, Pakistan.
[3]Department of Food Science and Technology, Jinnah University for Women Karachi, Pakistan
[4]Department of Pharmacy Practice, Jinnah University for Women Karachi, Pakistan

***For Correspondence**

Dr. Sidra Zubair
E-mail: phr.sidra@gmail.com, sidra.zubair@juw.edu.pk ; contact no. +92333-2213891

ABSTRACT

Essential oils are highly concentrated, volatile compounds extracted from various parts of plants, known for their unique aromas and potential therapeutic benefits. This review article explores the diverse applications, pharmacological activities, and current research on essential oils, with a focus on their antimicrobial, analgesic, and anti-inflammatory properties. The pharmacological activity of essential oils depends on their potency and route of administration. While oral ingestion is generally not recommended due to potential dangers, topical application with heat can enhance absorption. Essential oils have been linked to the stimulation of the limbic region of the brain, leading to relaxation and calmness.

Many essential oils have demonstrated analgesic (pain-relieving) effects, with clove oil being a well-known example. The main component responsible for this action is eugenol, a phenylpropanoid found in high concentrations in clove buds and leaves. Studies have shown the anti-nociceptive (pain-inhibiting) effects of eugenol and clove oil in animal models. Anti-inflammatory effects of essential oils are achieved by regulating the release of inflammatory mediators and inhibiting enzymes like cyclooxygenase. Essential oils from plants like Chamaecyparis obtusa have exhibited potent anti-oxidant and anti-inflammatory properties in in vitro and in vivo studies.

Essential oils also possess strong antimicrobial activities against a wide range of microorganisms, including bacteria, viruses, fungi, and protozoa. The antimicrobial effects are often attributed to the presence of terpenoids and other secondary metabolites. The minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) are used to evaluate the antimicrobial potency of essential oils. This review highlights the potential of essential oils as alternative therapies with fewer side effects compared to conventional medicines. However, further research is needed to fully understand the mechanisms of action, optimize therapeutic applications, and ensure safe and effective use of essential oils.

Keywords: Essential oils, Aromatherapy, Analgesic, Anti-inflammatory, Antimicrobial, Eugenol, Terpenoids.

INTRODUCTION

Essential oils are highly concentrated, volatile compounds extracted from various parts of plants, such as flowers, leaves, stems, and roots[(1)]. These oils are often referred to as the "essence" of the plant, as they capture the unique aroma and properties of the source material[(1)]. The term "essential" refers to the essence or quintessential nature of the oil, which is believed to represent the plant's unique characteristics.

Essential oils are typically obtained through physical methods like steam distillation, water distillation, or cold-pressing, rather than through chemical extraction[(2)]. These oils are highly concentrated and hydrophobic, meaning they are not soluble in water. They are known for their distinctive scents and are widely used in aromatherapy, personal care products, and various other applications[(3)].

The major constituents of essential oils can be divided into two main categories: hydrocarbons and oxygenated compounds[(4)]. Hydrocarbons include sesquiterpenes, monoterpenes, and diterpenes, while oxygenated compounds consist of aldehydes, esters, ketones, phenols, and other functional groups.

ISSN Online 3049-7647 © 2024 Elysian Sciences Publishing House

Aromatherapy, a form of alternative therapy, utilizes these essential oils for their potential therapeutic benefits. While the efficacy of essential oils in treating or curing diseases is still a subject of ongoing debate, their use has significantly increased in recent years. Many essential oils have demonstrated various pharmacological activities, such as analgesic (pain-relieving), anti-inflammatory, and antioxidant properties, depending on their specific chemical composition and the plant source[5, 6].

The following table provides an overview of the plant parts commonly used for the extraction of essential oils[3, 7]:

Table 1: overview of the plant parts commonly used for the extraction of essential oils

Bark	Berries	Flowers	Leaves	Resin	Rhizome	Roots	Seeds	Woods	Peel
Cassia	Allspice	Cannabis	Basil	Benzoin	Galangal	Valerian	Anise	Agarwood	Bergamot
Cinnamon	Juniper	Chamomile	Bay leaf	Copaiba	Ginger		Buchu	Camphor	Grapefruit
Sassafras		Clary sage	Buchu	Frankincense			Celery	Cedar	Lemon
		Clove	Cinnamon	Labdanum			Cumin	Rosewood	Lime
		Hops	Common sage	Myrrh			Flax	Sandalwood	Orange
		Hyssop	Eucalyptus				Nutmeg oil		Tangerine
		Jasmine	Guava						
		Lavender	Lemon grass						
		Manuka	Melaleuca						
		Marjoram	Oregano						
		Orange	Patchouli						
		Pelargonium (Scented geranium)	Peppermint						
		Plumeria	Pine						
		Rose	Rosemary						
		Ylang-ylang	Spearmint						
			Tea tree						
			Thyme						
			Tsuga						
			Wintergreen						

This review article aims to explore the diverse applications, potential benefits, and current research on the use of essential oils in various fields, including aromatherapy, personal care, and potential therapeutic applications.

Pharmacological Activity and Applications of Essential Oils

The pharmacological activity of essential oils depends on their potency and the route of administration into the body. Certain methods, such as topical application with the addition of heat, can increase the absorption of essential oils into the body. However, oral ingestion of essential oils is generally not recommended, as high concentrations can have dangerous effects, including burning sensations, salivation, and antispasmodic effects[6].

The use of essential oils has been linked to the stimulation of the limbic region of the brain, leading to relaxation and calmness. This is because the limbic

system is responsible for controlling various physiological functions, such as heart rate, breathing, and blood pressure. Alterations in the limbic system can directly or indirectly affect the state of mind and the body[8].

More than 90 types of essential oils are found in the world, and many have been extensively used for their beneficial activities. Some of the commonly used essential oils and their therapeutic applications include[9]:

Therapeutic Applications of Essential Oils

- Peppermint: Used to boost energy and aid digestion.
- Lavender: Used for stress relief.
- Sandalwood: Used to calm nerves and improve focus.
- Bergamot: Used to reduce stress and improve skin conditions like eczema.
- Rose: Used to improve mood and reduce anxiety.
- Chamomile: Used for improving mood and relaxation.
- Ylang-Ylang: Used to treat headaches, nausea, and skin conditions.
- Tea Tree: Used to fight infections and boost immunity.
- Jasmine: Used to help with depression, childbirth, and libido.
- Lemon: Used to aid digestion, improve mood, and relieve headaches.

Essential oils and aromatherapy have been suggested for the following common conditions[9, 10]:

- Depression and anxiety
- Headaches and migraines
- Sleeplessness or insomnia
- Analgesic (pain-relieving) effects
- Anti-inflammatory effects
- Antioxidant effects
- Antimicrobial and antibiotic properties
- Hormonal balance
- Boosting immunity and fighting infections
- Aiding digestion
- Increasing energy levels
- Improving skin and hair health
- Antimicrobial, Analgesic, and Anti-inflammatory Properties of Essential Oils

  This research article will focus on the antimicrobial, analgesic, and anti-inflammatory properties of essential oils. To understand these properties, it is important to identify the specific compounds responsible for these activities.

Sesquiterpenes and sesquiterpene alcohols are some of the compounds that possess anti-inflammatory properties[11]. Cineole, a terpene oxide commonly found in essential oils like eucalyptus, basil, cinnamon, and rosemary, has been shown to exhibit inhibitory effects on nociception (pain perception) receptors and reduce chemical-based inflammatory responses in experimental studies[11].

Other essential oil components, such as farnesene, limonene, pinene, aldehydes, and phenols, have demonstrated antibacterial, antiviral, or antifungal activities[12]. The effectiveness of these compounds is dependent on their concentration, dosage, and the overall potency of the essential oil.

Studies have shown that essential oils tend to be more effective at lower pH levels, lower temperatures, and in low-oxygen environments. Additionally, gram-positive microorganisms are generally more sensitive to essential oils compared to their gram-negative counterparts[13].

Specific essential oil components, such as thymol, carvacrol, and eugenol, have been found to have effective antibacterial properties, with minimum inhibitory concentrations (MICs) ranging from 0.05 to 5 microliters per milliliter in in vitro studies. In vitro studies have also demonstrated the antibacterial properties of essential oils against organisms like Escherichia coli, Listeria monocytogenes, Staphylococcus aureus, and Shigella dysenteriae at concentrations between 0.2 and 10 microliters per milliliter[14].

It is important to note that the composition and therapeutic properties of essential oils can be influenced by various factors, such as the season, climate, and the stage of plant maturity. These environmental factors can affect the concentration of the active compounds responsible for the antimicrobial, analgesic, and anti-inflammatory activities of the essential oils. Therefore, it is crucial to conduct studies that account for these variables to ensure accurate and reliable results[15].

Analgesic and Anti-Inflammatory Effects of Essential Oils

Analgesic effects refer to the ability of a substance, such as a drug or essential oil, to provide relief from pain by acting on the central and peripheral nervous systems. Analgesics differ from anesthetics, which can partially or completely eliminate sensation. Essential oils do not cause loss of sensation like anesthetics[16].

Clove oil, extracted from the herb Eugenia caryophyllata, is a well-known essential oil used for its analgesic effects, particularly for toothaches. The main component responsible for this action is eugenol, a phenylpropanoid that makes up 80-90% of clove buds and 82-88% of clove leaves. Eugenol has proven effective against a wide range of oral bacteria and viruses and is used in dentistry for dental caries, periodontal diseases, and root canal sealing[17].

Anti-inflammatory effects refer to the ability to reduce inflammation, which is a response to infectious agents, toxic substances, or physical injury. Inflammation involves the release of chemical mediators like histamine, cytokines, interleukins, and prostaglandins, leading to redness, swelling, pain, and heat. Inflammation can be acute, such as allergic reactions, or chronic, as in autoimmune diseases.

Many essential oils have demonstrated anti-inflammatory properties, including clove oil, lavender (Lavandula augustifolia Mill.), eucalyptus, and orange (Citrus sinensis). These oils can reduce inflammation by regulating the release of inflammatory cytokines and inhibiting enzymes like cyclooxygenase[16].




Figure 1: Structure of Some Essential oils

While conventional anti-inflammatory drugs like NSAIDs and corticosteroids are effective, their long-term use can lead to side effects, making essential oils an attractive alternative for some people

Antimicrobial Mechanisms of Essential Oils

Essential oils have demonstrated potent antimicrobial activities against a wide range of microorganisms, including bacteria, viruses, fungi, and protozoa. The antimicrobial effects of essential oils can be either biostatic, inhibiting the growth of microorganisms temporarily, or microbicidal, killing or eradicating the microorganisms completely[18].

Table 2: Antimicrobial Spectrum of Essential oil

Essential Oil	Plant Part	Chemical Composition	Antibacterial Activity Against Human Pathogens
Achillea fragrantissima	Aerial parts	Yomogi alcohol, 1,8-cineole, artemisia alcohol, thujone	Staphylococcus aureus, Staphylococcus epidermidis, Escherichia coli
Achillea ligustica	Aerial parts	Viridiflorol, terpinen-4-ol	Streptococcus mutans
Artemisia absinthium	Aerial parts	Myrcene, trans-thujone, trans-sabinyl acetate	E. coli, S. aureus, Staphylococcus epidermidis
Artemisia biennis	Aerial parts	(Z)-Beta-ocimene, (E)-beta-farnesene, acetylens, (Z)- and (E)-En-yn-dicycloethers	E. coli, S. aureus, S. epidermidis
Artemisia cana	Aerial parts	Santolina triene, alpha-pinene, camphene	E. coli, S. aureus, S. epidermidis
Artemisia dracunculus	Aerial parts	Methylchavicol, methyl eugenol, beta-phellandrene, terpinolene	E. coli, S. aureus, S. epidermidis, Brochothrix thermosphacta, Listeria innocua, L. monocytogenes, Pseudomonas putida, Shewanella putrefaciens
Artemisia longifolia	Aerial parts	Alpha-pinene, camphene, 1,8-cineole	E. coli, S. aureus, S. epidermidis
Artemisia frigida	Aerial parts	1,8-Cineole, methylchavicol, camphor	E. coli, S. aureus, S. epidermidis
Cinnamomum zeylanicum	Bark, leaves	Cinnamaldehyde	Enterobacteriaceae, S. aureus, Streptococcus pyogenes, S. pneumoniae, Enterococcus faecalis, E. faecium, Bacillus cereus, Acinetobacter lwoffii, Enterobacter aerogenes, E. coli, Klebsiella pneumoniae, Proteus mirabilis, P. aeruginosa, Salmonella typhimurium, Clostridium perfringens, Mycobacterium smegmatis

The antimicrobial activity of essential oils is often attributed to the presence of terpenoids and other secondary metabolites. Many essential oils derived from plants like thyme, clove, bay, and cinnamon have shown significant antimicrobial properties.

The antimicrobial activity of essential oils is typically evaluated by determining the minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC). These values can be assessed using the agar plate method, where the essential oil is placed on a filter paper and incubated with the target microorganism on an agar plate. However, it is important to note that different essential oils may have varying modes of action and may not be equally effective against all microorganisms.

Studies have revealed that some essential oils, such as sandalwood and vetiver, exhibit greater inhibitory effects on gram-positive bacteria compared to gram-negative bacteria. Conversely, other essential oils may be more effective against gram-negative strains. Additionally, certain essential oils may require specific conditions, such as temperature, pH, and oxygen levels, for optimal antimicrobial activity[8].

The antimicrobial mechanisms of essential oils involve disrupting the structural and functional integrity of the microbial cell membrane, leading to the leakage of intracellular substances and ultimately cell death. Essential oils can also interfere with the microbial efflux system, inhibiting the expulsion of antimicrobial agents from the cell.

Ongoing research is focused on exploring the antimicrobial potential of various essential oils and understanding their specific mechanisms of action against

ISSN Online 3049-7647 © 2024 Elysian Sciences Publishing House

different microorganisms. This knowledge can aid in developing novel antimicrobial agents and strategies to combat microbial infections while minimizing adverse effects associated with traditional antimicrobials[8].

The inhibitory effect of these oils was due to the occurrence of eugenol and cinnamaldehyde. Other major compounds were thymol, camphor and carvacrol.

Essential oils may disrupt the cell membrane of the targeted pathogens by increasing membrane permeability, inducing leakage of vital intracellular constituents, and interrupting the cellular metabolism and enzyme kinetics of the targeted pathogens. The present study reveals more information on in vitro research studies of essential oils; however, more efforts are required to conduct clinical trials in the future.

REFERENCES

1. 1. Hanif MA, Nisar S, Khan GS, Mushtaq Z, Zubair M. Essential oils. Essential oil research: trends in biosynthesis, analytics, industrial applications and biotechnological production. 2019:3-17.
2. Aziz ZA, Ahmad A, Setapar SHM, Karakucuk A, Azim MM, Lokhat D, et al. Essential oils: extraction techniques, pharmaceutical and therapeutic potential-a review. Current drug metabolism. 2018;19(13):1100-10.
3. Reddy DN. Essential oils extracted from medicinal plants and their applications. Natural Bio-Active Compounds: Volume 1: Production and Applications. 2019:237-83.
4. Mohammadhosseini M. Chemical composition of the essential oils and volatile fractions from flowers, stems and roots of Salvia multicaulis Vahl. by using MAHD, SFME and HS-SPME methods. Journal of essential oil bearing plants. 2015;18(6):1360-71.
5. Murti Y, Jain D, Semwal BC, Sıngh S, Janmeda P, Bhaskar P. Innovative methods for extraction of essential oils from medicinal plants. International Journal of Secondary Metabolite. 2023;10(2):190-230.
6. Dagli N, Dagli R, Mahmoud RS, Baroudi K. Essential oils, their therapeutic properties, and implication in dentistry: A review. Journal of International Society of Preventive and Community Dentistry. 2015;5(5):335-40.
7. Franz C, Novak J. Sources of essential oils. Handbook of essential oils: CRC Press; 2020. p. 41-83.
8. Chouhan S, Sharma K, Guleria S. Antimicrobial activity of some essential oils—present status and future perspectives. Medicines. 2017;4(3):58.
9. Pandey AK, Kumar P, Singh P, Tripathi NN, Bajpai VK. Essential oils: Sources of antimicrobials and food preservatives. Frontiers in microbiology. 2017;7:228506.
10. Baptista-Silva S, Borges S, Ramos OL, Pintado M, Sarmento B. The progress of essential oils as potential therapeutic agents: A review. Journal of Essential Oil Research. 2020;32(4):279-95.
11. Chan W-K, Tan LT-H, Chan K-G, Lee L-H, Goh B-H. Nerolidol: a sesquiterpene alcohol with multi-faceted pharmacological and biological activities. Molecules. 2016;21(5):529.
12. Tariq S, Wani S, Rasool W, Shafi K, Bhat MA, Prabhakar A, et al. A comprehensive review of the antibacterial, antifungal and antiviral potential of essential oils and their chemical constituents against drug-resistant microbial pathogens. Microbial pathogenesis. 2019;134:103580.
13. Oliva A, Costantini S, De Angelis M, Garzoli S, Božović M, Mascellino MT, et al. High potency of melaleuca alternifolia essential oil against multi-drug resistant gram-negative bacteria and methicillin-resistant Staphylococcus aureus. Molecules. 2018;23(10):2584.
14. Kachur K, Suntres Z. The antibacterial properties of phenolic isomers, carvacrol and thymol. Critical reviews in food science and nutrition. 2020;60(18):3042-53.
15. Marchese A, Orhan IE, Daglia M, Barbieri R, Di Lorenzo A, Nabavi SF, et al. Antibacterial and antifungal activities of thymol: A brief review of the literature. Food chemistry. 2016;210:402-14.
16. SILVA GL, Luft C, Lunardelli A, Amaral RH, MELO DA, Donadio MV, et al. Antioxidant, analgesic and anti-inflammatory effects of lavender essential oil. Anais da Academia Brasileira de Ciências. 2015;87:1397-408.
17. Khalilzadeh E, Hazrati R, Saiah GV. Effects of topical and systemic administration of Eugenia caryophyllata buds essential oil on corneal anesthesia and analgesia. Research in pharmaceutical sciences. 2016;11(4):293-302.

18. Swamy MK, Akhtar MS, Sinniah UR. Antimicrobial properties of plant essential oils against human pathogens and their mode of action: an updated review. Evidence-based complementary and alternative medicine. 2016;2016.

Received on:Dec 2023 Accepted on:March 2024 Published on Sep 2024

DOI:

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium provided the original work is properly cited.